WV BIRD RECORDS COMMITTEE REPORT FORM

Please Type or Print Legibly

Please Type or Print Legioly
Please Type or Print Legisly Number: (leave blank) Location: Beard WV, Pocahontas Co. Date: Dec. 14, 2016 Time: 11:30 a.m. Bird Seen: Dec. 10, 2016 To: Dec. 14, 2016 Age/Sex: Unknawn Description: In lieu of physical material substantiating a record (e.g., specimen, diagnostic photograph, vocal recordings) a complete written description can serve as verification for the biological record. Please be as thorough as your field notes allow. Do not feel constrained by the space provided. Additional supporting descriptions/illustrations may be attached. Note that photographs or recordings WILL NOT BE RETURNED but become part of the permanent record for the observation. Structure: (Overall size and shape plus details on the head, feet, tail, etc.) "Medium - sized black bird "Smaller than a common grackle. Larger than a cowbird." "Smaller than a common grackle and perhaps even a little shorter than one might expect for a bird of that size. "Black tegs and beak"
· Black regs and sour
Plumage: (e.g. markings, coloration, molt) 610554 black with a more dramatically irridescent head and neck. No brown observed at all.
Description of any vocalizations, if heard: None
Description of behavior: Not seen on the feeders above the ground, Seen feeding on the ground on (presumed) bits of seeds under the blackoil the ground on (presumed) bits of seeds under the safflower sun flower feeder and to a lesser extent on the safflower feeder. Also foraged in the leef litter where it walked feeder. Also foraged in the leef litter where it walked about, tossing leaves. Gait not that of previously about, tossing leaves. Gait not that of previously observed common grackle or combined. Not like a rust blackbird (seen previously in a nearby wetland area

Sheaf litter re allowed to accomplate and a couple of brush Piles provide additional shelter. Habitat: (General and Specific) Walley with open pasture land bordered
Sheaf litter re allowed to mal shelter,
Habitat: (General and Specific) Maintainous river valley with open pasture land bordered
M Lack - Front
by mature mixed hardwood forest. The novel elected by mature mixed hardwood forest. The novel elected with a areas where observed are on the forest edge with a areas where observed are on the forest edge with a areas where observed and adjacent pastureland
areas where observa small provid and adjacent passicione
areas where observed are on the forest eage form to areas where observed are on the forest eage format pastureland yard of 3-4 acres, a small point and adjacent pastureland with horses. The feeder area is roughly by acre scattered with horses. The feeder area is roughly by acres cattered
with horses. The factor area is roughly with horses. The factor and includes stands forbirds and trees around the hase shrubs and includes stands forbirds and trees
To the set of observation: (Include details of distance to bird, optics employed, righting
First observed the morning of Dee 10 armenises at Saffliner feeder. Appeared intermittently in the morning saffliner feeder. Appeared intermittently in the morning saffliner feeder. Appeared intermittently in the morning
house about 20-30 feet away. Queric's made to others
and report to e Bird filed the ty, wash to fix a scope
First observed the morning safflower feeder. Appeared intermi Hently in the morning safflower feeder. Appeared intermi Hently in the morning hours about 20-30 feet away. (Queries made to others hours about 20-30 feet away. (Queries made to others and report to e Bird filed Dec. 14, 2016.) Viable to fix a scope Used Swarovski. EL 8.5 x 42 binoculars. (Viable to fix a scope Used Swarovski. EL 8.5 x 42 binoculars.) Pictures taken with Apple iphone
Used Swarovski EL 8.5 x 42 binoculars. Unable to tix a scope of the bird.) Pictures taken with Apple iphone due to transient nature of the bird.) Pictures taken with Apple iphone
Previous experience with this species: Limited, May have seen in Arizona 410 years ago.
· ·
Experiences with similarly appearing species which are eliminated by your description: (Please be specific)
Often see Brown neader conting Within a mile or so
bird was different from the others see page 3-
- See parge -
Other observers: (Names, Addresses, E-mail please)
If the others agree with your identification they can sign this form here. If they care to provide additional details they can submit their own report.
7
Additional observer signature:
Additional observer signature:
Additional cook for organisms.
Additional observer signature:

Books, illustrations and advice consulted, and especially how did these influence this description: - Shared pic on Falebook in "What's this Bird?" and "Mountain state Birders" with mixed comments for Rosty Brewers ! State Birders" with mixed comments for Rosty Brewers! Common Grackle. Made decision based on my experience with hirds on the beard areas	
Books, illustrations and advice constituted, and expenses in "What's this Bird: alle revers!	
- Shaved pic " with mixed comments for hesightena with	
State 1311 cers Grackle. Made decision based on my com	
Common Grackle. Made areas birds in the beard areas (Goide to Brids) Mitch Waite Group i Phone app "iBird Pro" (Goide to Brids of North America)	
; Phone app "iBird for Burds of North America)	
- All agon is less de	
5 C/O/2 Court write this description?	
How long after observing this bird did you first write this destribution. Dec. 14, 2016 report to eBird	
Name:	
Address:	
Date:	
Signature:	

Mail to: WV Bird Records Committee
Attn: Wil Hershberger
170 Stallion Ct.
Hedgesville, WV 25427
Or E-mail to:
wilhershberger@mac.com



Please use the space below to add any appropriate details you feel were missing from the above form or to expand on any answers whose allotted space was insufficient for a complete response: Experiences (cont.) Not a Common gracke: True, it had a yellow eye-but the bird was smaller and had a "two-toned" Opinions appearance, Did not have a characteristic Grackle appearance, Did not have a characteristic Grackle tail and foraged differently in that it was more tail and foraged differently in that it was more deliberate, had a more measured goit, and overall seemed slower. Not a Rusty blackbird (also seen in a wetland area " a mil & eway previously). Did not have any brownstant "usty" appearance. Not a Brun-headed combird becowse of plumage and behavior. Also, the bird in question was larger. O Capy of Dec 14 e Bird report. (Note additional information was provided to Mike Slaven.)

Two pics - enlarged. [1 - 11-1-1 --1+ 3 March 11, 2017 eBird report of additional sighting Beard location is near Hillsboro WV. Location of sighting is our home "Ravens' Roost" @ 18 Lang. -80.2292. Addendum .

3/16/2017

Submit Observations

Explore Data

My eBird

Help

Sign In or Register

Language

Older

All Checklists

Newer

Location

Beard WV, Pocahontas County, West Virginia, US (Map)

Date and Effort Wed Dec 14, 2016 11:30 AM

Protocol:

Traveling

Party Size:

2

Duration:

3 hour(s), 15 minute(s)

Distance:

3.0 mile(s)

Observers:

Species

29 species total

Hide Media

- 8 Canada Goose
- 25 Mallard
 - 1 Northern Harrier

"Brown" female or juvenile hunting in the prison pond field.

4 Wilson's Snipe

Slumbering and feeding in the eastern shallows of the prison pond.

- 6 Mourning Dove
- 1 Belted Kingfisher

at Joel's pond

Age & Sex		Juvenile	Immature	Adult	Age Unknown
-					Will report to the first that the first the fi
	Male				
	Female				
	Sex Unknown				

- 1 Red-bellied Woodpecker
- 1 Yellow-bellied Sapsucker
- 3 Downy Woodpecker

- 1 Northern Flicker
- 1 Pileated Woodpecker
- 1 American Kestrel

Hunting in the area of the barn across from the Shaw's.

1 Eastern Phoebe

At the prison pond.

- 4 Blue Jay
- 9 American Crow
- 1 Common Raven
- 5 Black-capped Chickadee
- 8 Tufted Titmouse
- 1 White-breasted Nuthatch
- 2 Carolina Wren
- 1 Golden-crowned Kinglet
- 1 Northern Mockingbird
- 20 Dark-eyed Junco
- 1 White-throated Sparrow
- 4 Song Sparrow
- 2 Northern Cardinal

1 Brewer's Blackbird

Sighted in the vicinity of house feeders (for at least 4 days) feeding on the ground and in the leaf litter where it walked about, tossing leaves. Ruled out Common Grackle because of smaller size and short tail. Ruled out Rusty Blackbirds (also seen in the Beard area - but not at the house feeders in the past) because of absence of brown/tan/"rusty" appearance. Showed pic on Facebook in "What's this Bird?" and "Mountain State Birders" with mixed comments for Rusty/Brewers/Common Grackle. Made decision based on my experience with birds in the Beard area. Precise location in Longitude/Latitude of the bird can be provided.



Macaulay Library

The yellow eye is clearly visible in the individual although not as clear in the attached picture. The dark-iridescent "hood" is also evident in the individual.


2 Purple Finch

20 American Goldfinch

Are you submitting a complete checklist of the birds you were able to identify?

Yes

© Cornell Lab of Omithology | Contact | FAQ





March 11, 2017

From:

ebird-checklist@cornell.edu

Sent:

Saturday, March 11, 2017 9:01 PM

To:

Subject:

eBird Report - Beard WV, Mar 11, 2017

Beard WV, Pocahontas, West Virginia, US

Mar 11, 2017 4:50 PM - 6:30 PM

Protocol: Traveling

2.5 mile(s)

Comments: Sunny, no wind, late in the day, low 30s. Walked from Ravens' Roost to the prison pond and return.

22 species

Canada Goose 17

Wood Duck 11 On the prison pond. 5 females/6 males

Black Vulture 2

Turkey Vulture 30 Vultures were gathered around two cow carcasses placed in Hanley's "cow cemetery."

Mourning Dove 3

Red-bellied Woodpecker 1

Downy Woodpecker 1

American Crow 3

Common Raven 2

Black-capped Chickadee 2

Tufted Titmouse 2

White-breasted Nuthatch 2

American Robin 35 Saw two large flocks. Could have been the same one of 30+ individuals.

European Starling 5

Fox Sparrow 5 Observed together in close proximity in the brushy area about MP 39 before the prison pond.

Dark-eved Junco 4

White-throated Sparrow 1

Song Sparrow 4

Swamp Sparrow 1

Northern Cardinal 1

Red-winged Blackbird 30

Observed in a horse paddock near Beard (exact location can be provided). First sighted among a Brewer's Blackbird 1 mixed flock of 5 European Starlings and two dozen Red-winged blackbirds all grouped together. A flock of about 30 robins were nearby, too. Dramatically distinguished from the others by it's size, behavior, consistent solid iridescent black, yellow eyes, and a non-grackle tail and smaller size than a grackle. The individual then left the mixed flock and roamed in the grass near the scattered robin flock giving me a better look. It's behavior gave another distinction in that it's manner of foraging varied from the other noted birds.

I may have seen a second individual near the prison pond. Heard his call note, but was unable to get a recording. In December, I reported a Brewer's Blackbird at our feeders at our house across the road from the horse paddock noted in the current report. I am in the process of preparing a report for the Brooks Bird Club record committee on that observation. The bird I sighted today looked just like the bird reported in the Dec sighting.

View this checklist online at

This report was generated automatically by eBird v3 (http://ebird.org)