

Hairy Woodpecker (*Picoides villosus*)

WINTER: Uncommon to Fairly Common Resident. See Remarks section for CBC data.

SUMMER: Uncommon to Fairly Common Resident. See Remarks section.

MIGRATION:

Spring: Uncommon to Fairly Common Resident.

Fall: Uncommon to Fairly Common Resident.

REMARKS:

Singing Male Census (Eddy 2003): Cabwaylingo State Forest, Wayne Co., WV (2002): Twelvepole Creek: dominant canopy: yellow poplar, sugar maple, sycamore (elevation, 244 m at center line): +.

Breeding Bird Survey (Bullard 2003): Cabwaylingo State Forest, Wayne Co., WV (2002): 135 Stops (7 surveys), 64 Mile. Number of Stops Recorded: 3. Total Birds: 3.

Singing Male Census (Koch and Hurley 1972): Beech Fork Lake, Wayne Co., WV (1969): mature mesophytic forest (elevation, 198 m to 244 m): 8 males/100 ha.

CHRISTMAS BIRD COUNT DATA

Huntington, WV (1940 to 2001): Recorded on 100% of Counts. Total Birds: 462. Range: 1 to 37.

Mason County, WV (1953 to 1976): Recorded on 94% of Counts. Total Birds: 55. Range: 1 to 13.

Ona, WV (1962 to 2001): Recorded on 100% of Counts. Total Birds: 236. Range: 1 to 14.

Huntington, WV (1940 to 2001): Number of Counts: 61. Birds/Party Hr.: <1.0. Party Hr./Bird: 6.5.

Mason County, WV (1953 to 1976): Number of Counts: 18. Birds/Party Hr.: <1.0. Party Hr./Bird: 6.4.

Ona, WV (1962 to 2001): Number of Counts: 40. Birds/Party Hr.: <1.0. Party Hr./Bird: 7.7.