Red-shouldered Hawk (Buteo lineatus)

WINTER: Fairly Common Resident. See Remarks section for CBC data.

SUMMER: Fairly Common Resident.

MIGRATION:

Spring: Migration status uncertain due to insufficient data.

Fall: Uncommon Migrant at best. Data for migration dates lacking. Smith (1996) indicates possible migrants in October.

REMARKS:

Breeding Bird Survey (Bullard 2003): Cabwaylingo State Forest, Wayne Co., WV (2002): 135 Stops (7 surveys), 64 Mile. Number of Stops Recorded: 1. Total Birds: 1.

Breeding Bird Survey (Koch and Hurley 1972): Beech Fork Lake, Wayne Co., WV (29 May 1969): 50 Stops, 25 mile. Number of Stops Recorded: 1. Total Birds: 2.

Breeding Bird Survey (Koch 1974): McClintic WMA, Mason Co., WV (29 May 1971): 50 Stops, 12.25 Mile. Number of Stops Recorded: 1. Total Birds: 1.

CHRISTMAS BIRD COUNT DATA

Huntington, WV (1940 to 2001): Recorded on 75% of Counts. Total Birds: 200. Range: 1 to 18.

Mason County, WV (1953 to 1976): Recorded on 100% of Counts. Total Birds: 37. Range: 1 to 4.

Ona, WV (1962 to 2001): Recorded on 98% of Counts. Total Birds: 199. Range: 1 to 13.

Huntington, WV (1940 to 2001): Number of Counts: 61. Birds/Party Hr.: <1.0. Party Hr./Bird: 15.1.

Mason County, WV (1940 to 2001): Number of Counts: 01. Birds/Party Hr.: <1.0. Party Hr./Bird: 9.5. Ona, WV (1962 to 2001): Number of Counts: 40. Birds/Party Hr.: <1.0. Party Hr./Bird: 9.1.