## Sharp-shinned Hawk (Accipiter striatus)

WINTER: Uncommon Resident. See Remarks section for CBC data.

**SUMMER:** Uncommon Resident.

## **MIGRATION:**

Spring: Uncommon Migrant. Kiff et al. (1986) indicates migration occurs in April and May.

Fall: Uncommon Migrant. Slack (1987) indicates migrants occur in October.

## **REMARKS:**

See Remarks section for Cooper's Hawk.

Singing Male Census (Eddy 2003): Cabwaylingo State Forest, Wayne Co., WV (2002): Tick Ridge: dominant canopy: yellow poplar, black cherry, sugar maple (elevation, 350 m to 378 m): +

Banding Data: One banded (LK) on 31 December 1963 at Ona, Cabell Co., WV was recovered in April 1964 in North Carolina (Kiff et al. 1986).

## **CHRISTMAS BIRD COUNT DATA**

Huntington, WV (1940 to 2001): Recorded on 52% of Counts. Total Birds: 61. Range: 1 to 5. Mason County, WV (1953 to 1976): Recorded on 22% of Counts. Total Birds: 5. Range: 1 to 2. Ona, WV (1962 to 2001): Recorded on 75% of Counts. Total Birds: 71. Range: 1 to 10.

Huntington, WV (1940 to 2001): Number of Counts: 61. Birds/Party Hr.: <1.0. Party Hr./Bird: 49.5. Mason County, WV (1953 to 1976): Number of Counts: 18. Birds/Party Hr.: <1.0. Party Hr./Bird: 70.6. Ona, WV (1962 to 2001): Number of Counts: 40. Birds/Party Hr.: <1.0. Party Hr./Bird: 25.6.